

## Report on Oribatid Mites in Eco-Friendly Agriculture with a Description of Third New Species of the Genus *Cosmogalumna* from the Litter of Coconut Palm Tree on Ishigaki Island, in Southern Japan

YOSHIO NAKAMURA\* and TOKUKO FUJIKAWA\*\*

### Abstract

Oribatids are subdivided taxonomically into Macropylina, Apterogastrina and Pterogasterina. In the field of eco-friendly agriculture, the number of Pterogasterina species was clearly larger than in the field with tillage management or chemical fertilizer application. This indicates that an increase in the number of Pterogasterina species is a bio-indicator of eco-friendly agriculture. A new species of the Pterogasterina, *Cosmogalumna hiroyoshii*, was collected from coconut palm tree litter on Ishigaki Island, in southern Japan. Characteristic features of the new species include thick rostral setae which are not extending in front of the anterior rostral margin; a bimucronate rostrum; setae *le* inserted lateral to the carina L between L and L; the polygonal structure on the prodorsal, notogastral, pteromorph surfaces, and ventral plate; and on the alveoli of dorsal setae, granules on and around each area porosa.

**Key words:** Pterogasterina, *Cosmogalumna hiroyoshii* sp. nov., Oribatid mites, Ishigaki Is., Eco-friendly Agriculture

Soil animals are divided into four categories by the method in which they are collected: macrofauna, dry mesofauna, wet mesofauna and microfauna (Nakamura, 2000). Together with collembola, oribatid mites are the main component in the dry mesofauna of cropfield soil (Nakamura, 1988; Nakamura *et al.*, 2000). Oribatid mites are about 0.1–1.6 mm in body length, and have a dark brown and chitinized body. About 9,000 species are known world-wide, and 900 species are found in Japan (Balogh, 1972; Fujikawa, 1991; Fujikawa *et al.*, 1993). In a broad sense, they are saprophagous (Wallwork, 1976) and are often called ‘moss mites’. But many of them feed on fungi (Matsuzaki and Itakura, 1991; Nakamura *et al.*, 1991, Nakamura, 1993). A recently discovered species, *Schelorbitates azumaensis* ENAMI, NAKAMURA *et* KATSUMATA, 1996, from Japanese cropfields, actively ate *Rhizoctonia solani* KÜHN, a plant pathogenic fungus, and prevented root rot of radish under laboratory conditions (Enami and Nakamura, 1996). This indicates that oribatid mites possess great potential as agents for the biological control of soil-borne crop root diseases.

Recently, eco-friendly agriculture has been emphasized. Based on the opinion that eco-friendly agriculture should also be friendly to soil animals (cf. Nakamura, 1998), studies on the effect of various agricultural practices on

soil animals have been carried out. For instance, soil animals were more numerous and complicated in no-tillage cultivation than in tillage cultivation (Nakamura, 1988), or in organic farming than in chemical-intensive farming (Fujikawa, 1976; Nakamura *et al.*, 2000). Organic mulch of crop residues and chopped weeds help create a healthy environment for soil animals, especially enchytraeids and earthworms (Nakamura *et al.*, 2003). Trials involving the artificial introduction of soil animals into a crop field proved that a certain species could colonize the crop fields when the introduction was carried out together with forest litter (oribatid mites by Fujikawa, 1988; macro animals by Nakamura, 1998). This suggests that organic materials transferred into crop fields provide a medium for settlement of soil animals, in addition to supplying organic fertilizer for the crops. In tropical regions, coconut palm litter is thought to be an important mulch material.

Since oribatids are subdivided taxonomically into two types: the Lower (Primitive) Oribatei (=Macropylina; ex. *Eniochthonius minutissimus* BERLESE, 1904, Fig. 1A) and the Higher Oribatei (=Brachypylylina) of the Apterogasterina (ex. *Oppiella nova* (OUDEMANS, 1902), Fig. 1B) and Pterogasterina (Balogh, 1972), the species complex plays a bioindicator in agroecosystems (Behan-Pelletier, 1999; Enami, 2000). In the natural forest of Japan, the number of Apterogasterina species was numerous (Fujikawa, 2004). In the crop fields, on the other hand, the number of Pterogasterina species was numerous (Nakamura, 1989; Enami, 2000). And the number of Pterogasterina species was clearly larger in fields where no-tillage was practiced (Nakamura, 1989) or organic materials were applied to the surface soil (Fujikawa, 1988; Nakamura, 1998), as

Received 8 March 2004.

Accepted 17 May 2004.

\*Laboratory of Crop Sciences, Faculty of Agriculture, Ehime University, 3-5-7 Tarumi, Matsuyama city, Ehime, 790-8566, Japan

\*\*Yokogawara 1375, Shigenobu town, Ehime, 791-0203, Japan




Fig. 1. Species of the Lower (Primitive) and the Higher oribatids. A: *Eniochthonius minutissimus* (BERLESE, 1904) of the Lower oribatids, B: *Oppiella nova* (OUDEMANS, 1902) of Apterogasterina, the Higher, C: *Transoribates agricola* NAKAMURA et AOKI, 1993 of Pterogasterina, the Higher, and D: The new species of Pterogasterina, the Higher. Black arrows show pteromorphae.




Fig. 2. *Cosmogalumna hiroyoshii* sp. nov. NSMT-Ac 11738 and 11739. A: Dorsal view, B: Prodersum, C: Pteromorphae and D: Anal region.

compared with tillage management or chemical fertilizer application. Furthermore, all of the dominant species belong to Pterogasterina; *Eporibatula sakamorii* (AOKI, 1970) in the cropped heavy clay soil of Nayoro Nature Farm, the northern district (Fujikawa, 1988), and *Transoribates agricola* NAKAMURA et AOKI, 1989 (Fig. 1 C) in the cropped andosol of National Inst. Agro-Environmental Science, Tsukuba, the central district (Nakamura, 1989), respectively. In the cropped andosol of National Tohoku Agric. Exp. Stn., Fukushima, the northern district, the number of individuals of *Sche-loribates azumaensis* of Pterogasterina was enormous (Nakamura, 1998). These results suggest that an increase in the number of Pterogasterina species is a bio-indicator of eco-friendly agriculture.

The new species being reported here also belongs to the genus *Cosmogalumna*, one of the members of the Pterogasterina group, which have well developed pteromorphae and area porosae (Fig. 1D). The pteromorphae of this species are especially clear and thick, with a large light colored polygonal reticulated pattern (Fig. 2A & B) which is covering whole body. The question is its function: do the pteromorphae protect against enemies and desiccation by covering the body, and/or against an increase of body temperature by fanning, or do they help it to fly for wind dispersion as Collembola suggested by Farrow and Greenslade (1992). The various species with such a large polygonal network are collected from tropical or subtropical areas. Other questions are whether the network is helpful for heat radiation, or for collecting raindrop by expansion of the body surface area. The present new species, *Cosmogalumna hiroyoshii*, was collected from coconut palm tree litter on Ishigaki Island. In tropical district including Ishigaki Island, the palm tree litter is thought to be the mulch and compost materials.

In 1988 Aoki defined the genus *Cosmogalumna*, and described a new, and type, species, *Cosmogalumna ornata*, from Nakanoshima Island in the Tokara Islands (Aoki, 1988). So far as the authors know, only one species has been added to this genus. The second species, *C. imperfecta* AOKI et HU, 1993 was collected from Yunnan Province in southern China (Aoki and Hu, 1993). The species described in this report was also collected from a subtropical region, Ishigaki Island, and is the third member of the genus. The notation follows that of Hammen (1980).

*Cosmogalumna hiroyoshii* sp. nov.

[Japanese name: Hiroyoshi-kazarifurisodedani]

**Measurements** (2 exs.) **and color:** Body length, 336  $\mu\text{m}$ ; wide, 257  $\mu\text{m}$ . Light yellowish brown.

**Body surface:** The whole integument on prodorsum, notogaster, pteromorphae and ventral plate with fine granules and distinct light colored polygonal network. Lamellar region of prodorsum, on and around area porosae, genital and anal plates densely covered by irregular dark colored granules. Femora with striate pattern of granules.

**Prodorsum:** According to depressed specimen, rostrum bimucronate; mucrones convergent, forming a trapezoid concavity (Fig. 3C). A large elliptical light area of weak chitinisation present just behind the concavity. Rostral setae (*ro*) thick, originating ventrally. Lamellar setae (*le*) shorter than *ro*, inserted between carinae L and L', just lateral to carinae L. Carinae extending for about half length of prodorsum. Interlamellar setae (*in*) and ex-oboithridial setae (*ex*) located at the same level. Setae *le*, *in* and *ex* short, setiform and slightly roughened. Sensilli (*ss*) composed of a thin stem and an expanded head which bears bristles; *ss* extending at the level of antero-margin of carinae (Fig. 3A). Relative lengths:  $ss > ro > le > ex > in$  (Fig. 3D).

**Notogaster:** Circular in form with a small indentation at the medial portion of the posterior margin which is not seen in depressed specimen. Length as long as, or slightly longer than width. Pteromorphae extending for a short distance in front of carinae L and S on prodorsum; furrow (*st* after Grandjean, 1966) remarkable (Fig. 3E). Five pairs of area porosae present; the relative sizes:  $Ad > Aa > A_1 > A_3 > A_2$  (Figs. 3F-H). Ten pairs of alveoli of dorsal setae present. Lyrifissures *ia*, *im*, *ip* and *ips* discernible; *im* located at inverse apomedial line and antero-laterally to  $A_1$ .

**Ventral region:** Genito-anal setae: (6 or 5-1-2-3); genital setae variable in number; all setae short and glabrous (Fig. 4D). Relative lengths:  $g_1 > g_2$  to  $g_6 > ad_1 > an > ag \doteq ad_2, ad_3$ . Lyrifissures *iad* situated parallel to the lateral side of anal aperture and almost mid-distance between the levels of two anal setae (Fig. 4E). Adanal setae  $ad_3$  inserted lateral to *iad*. Epimeral setae *1a*, *3b* and *4a* present; relative lengths:  $3b \doteq 3 \times 1a$ ;  $1a \doteq 4a$ . Alveoli *4b* present (Fig. 4B). Subcapitulum diarthric, bearing each one pair of anterior (*a*), medial (*m*) and posterior (*h*) infracapitular setae (Fig. 4A); all setae smooth; relative lengths:  $a \doteq h > m$ . Genito-anal, epimeral and infracapitular setae inserted at the end of a dark colored inner capsule.

**Legs:** All legs heterotridactylous; median claw thicker than lateral ones, with a dent ventrally at the middle portion. Leg chaetotaxy including famulus, but excluding solenidia, I (1-4-3-4-20); II (1-4-3-4-15); III (2-2-1-3-15); IV (1-2-2-3-11). Solenidiotaxy, I (1-2-2); II (1-1-2); III (1-1-0); IV (0-1-0). On tarsus I, solenidion  $\omega_1$  bacilliform, coupled with  $ft''$ ; solenidion  $\omega_2$  setiform, inserted posterior-laterally to  $\omega_1$ ; famulus  $\epsilon$  setiform, shorter than  $ft''$ , inserted anterior to and between solenidia (Fig. 5A). All solenidia on leg II bacilliform (Fig. 5D), but solenidia on tibia I, genu I (Fig. 5B), tibia III (Fig. 5F), genu III (Fig. 5G) and tibia IV (Fig. 5H) setiform. Trochanter III with  $v'$ , of long setiform bearing barbs unilaterally, and  $l'$  of remarkable like a trigonal pyramid (black arrow in Fig. 5G).

**Material examined:** Holotype (NSMT-Ac 11738) from litter of coconut palm tree (*Satakentia liukuensis* MOORE, 1969, Family Palmae) in Ishigaki Island, Okinawa Pref., Oct. 25, 2000, H. Shiraiishi; 1 paratype (NSMT-Ac




Fig. 3. *Cosmogalumna hiroyoshii* sp. nov. A & B: NSMT-Ac 11738; C to H: NSMT-Ac 11739. A: Dorsal view, B: Ventral view, C: Rostrum, D: Right lamellar region, E: Furrow on pteromorph, and F to H: Area porosal region. *ro*: Rostral seta; *le*: Lamellar seta; *in*: Interlamellar seta; *ex*: Exobothridial seta; *ss*: Sensillus; *ta*, *te*, *im*, *ms*, *r1*, *r3*, *p1* and *p2*: Dorsal setae; *Aa*, *A1*, *A2* and *A3*: Area porosae; *gla*: Glanular opening; *L*: Lamellar line; *S*: Sublamellar line; *st*: Furrow. Black arrows in Figs. E to H direct to rostrum.




Fig. 4. *Cosmogalumna hiroyoshii* sp. nov. NSMT-Ac 11739. A: Infracapitular region, B: Epimeral region, C: Coxal region of leg I to IV, D: Genital region, E: Anal region. *a*, *m* and *h*: Anterior, medial and posterior infracapitular setae, respectively; *an*: Anal seta; *ad*: Adanal seta; *ag*: Aggenital seta; *g*: Genital seta; *1a*, *3b* and *4a*: Epimeral setae; *4b*: Alveoli; *iad*: Adanal lyrifissure.




Fig. 5. *Cosmogalumna hiroyoshii* sp. nov. NSMT-Ac 11739. A: Right tarsus I, B: Right tibia I and genu I, C: Left femur I and a part of trochanter I, D: Left leg II, E: Right tarsus I, F: Right tibia III, G: Right genu III to trochanter III, H: Right tarsus IV and tibia IV, I: Left genu IV, J: Right femur IV and trochanter IV.

Table 1. Comparison in some features among three species of the genus *Cosmogalumna*

	<i>C. ornata</i> AOKI, 1988	<i>C. imperfecta</i> AOKI et HU, 1993	<i>C. hiroyoshi</i> sp. nov.
Ro	rounded	rounded	bimucronate
ro	fine; extending in front of anterior rostral margin	extending in front of anterior rostral margin	thick; not extending in front of anterior rostral margin; not seen from above
<i>le</i>	< <i>ro</i> ; between L and S	≐ <i>ro</i> ; lateral to L	< <i>ro</i> ; lateral to L
PN	notogaster, Ge-An	notogaster, Ge-An	prodorsum, notogaster ptero- morphae, ventral plate
BL	317(330)342µm	326–365µm	336µm
BW	256(264)270µm	276–287µm	257µm
Coll.	Mt. Mitake, Nakanoshima Is. of Tokara Islands	natural forest Xishuangbanna of Yunnan in China	coconut palm tree in Ishigaki Is., Okinawa prefecture

Ro: Rostrum; *ro*: Rostral seta; *le*: Lamellar seta; L: Lamellar line; S: Sublamellar line; PN: Polygonal network; Ge-An: Ventral surface between genital and anal apertures; BL: Body length; BW: Body width; Coll.: Place where samples were collected.

Table 2. Comparison in a few features among species of different genera with body surface covered by a reticulated pattern

	<i>Trachygalumna bisulcata</i> BALOGH, 1960	<i>Flagellozetes porosus</i> BALOGH, 1970	<i>Cosmogalumna</i> spp.
<i>le</i>	between L and L	between L and S	between L and S, or L and L
ds	short, partly absent	all long, flagellate	alveoli
Ap	punctiform	developed	developed
Gr	absent	absent	present
PN	notogaster, Ge-An pteromorphae	notogaster, prodorsum, pteromorphae	notogaster, Ge-An (prodorsum, pteromor- phae, notogaster & ventral plate of the new species)
BL	250µm	529µm	317–365µm
BW	202µm	329µm	256–287µm
Coll.	Africa	Ceylon	Tokara Is., Ishigaki Is., China

*le*: Lamellar seta; ds: Dorsal seta; Ap: Area porosae; Gr: An aggregation of granules on and around each area porosa; PN: Polygonal network; Ge-An: Ventral surface between genital and anal apertures; BL: Body length; BW: Body width; Coll.: Place where samples were collected.

11739): the same data as holotype. Types are deposited in the National Science Museum, Tokyo.

**Remarks:** The new species is conspicuous by the body surface covered by a network-like pattern. Surface structure of this kind is seen in some known species, for examples, *Cosmogalumna ornata* AOKI, 1988, *C. imperfecta* AOKI et HU, 1993, *Flagellozetes porosus* BALOGH, 1970 and *Trachygalumna bisulcata* BALOGH, 1960. However, the present species differs from *T. bisulcata* by having developed area porosae and polygonal structure on prodorsal surface, from *F. porosus* by having alveoli of dorsal setae, granules on and around each area porosa, and lamellar setae inserted between L and L, from *C. ornata* by having polygonal structure on prodorsal and pteromorph surface, thick rostral setae not extending in front of anterior rostral margin, bimucronate rostrum, and setae *le* inserted lateral to carinae between L and L, and from *C. imperfecta* by having polygonal structure on prodorsal and pteromorph surface, bimucronate rostrum, and rostral setae longer than lamellar setae, not extending in

front of anterior rostral margin, as summarized in Tables 1 and 2.

**Ethymology:** The new species was named in honor of the late Mr. Hiroyoshi SHIRAISHI (1968–2002), a researcher at the Tohoku National Agricultural Experiment Station. Mr. H. SHIRAISHI, while working at the Ministry of Agriculture, Forestry, and Fisheries of Japan, studied soil animals, especially oribatid mites (Shiraishi and Aoki, 1994) and springtails (Collembola) (Shiraishi *et al.*, 2003), as the agents for biological control of soil-borne crop root diseases. This new species was obtained from his samples during a survey for collembolan fauna in the subtropical Ishigaki Island.

## References

- Aoki, J. (1988) Oribatid mites (Acari: Oribatids) from the Tokara Islands, Southern Japan II. Bull. Biogeogr. Soc. Japan, 43(6): 31–33.
- Aoki, J. & S. Hu (1993) Oribatid mites from tropical forests of Yunnan Province in China II. Families Galumnidae and Galumnellidae. Zool. Sci., 10: 835–848.
- Balogh, J. (1960) Oribates (Acari) nouveaux d'Angola et du Congo Belge (2<sup>ème</sup> Série). Publ. cult. Co. Diam. Ang. Lisboa, 51: 14–40.
- Balogh, J. (1970) New oribatids (Acari) from Ceylon. The scientific results of the Hungarian Soil Zoological Expeditions. Opusc. Zool. Budapest, 10: 33–67.
- Balogh, J. (1972) The oribatid mites genera of the world. Akademiai, Budapest, 188pp, 72 plates.
- Behan-Pelletier, V. M. (1999) Oribatid mite biodiversity in agroecosystems: role for bioindication. Agr. Ecosystems Environ., 74: 411–423.
- Enami, Y. (2000) Oribatid mites as decomposers and their use as bioindicators of upland soil management. Farming Japan, 34(5): 16–21.
- Enami, Y. and Y. Nakamura (1996) Influence of *Schelorbitates azumaensis* (Acari: Oribatida) on *Rhizoctonia solani*, the cause of radish root rot. Pedobiologia, 40: 251–254.
- Farrow, R. A. and P. Greenslade (1992) A vertical migration of Collembola. The Entomologist, 111: 38–45.
- Fujikawa, T. (1976) Study on oribatid mites in nature farming and habitual farming fields. Edaphologia, 15: 1–11. (In Japanese.)
- Fujikawa, T. (1988) Fluctuation of oribatid mites in Nayoro nature farming field during ten years. Edaphologia, 39: 29–37.
- Fujikawa, T. (1991) List of oribatid families and genera of the World. Edaphologia, 46: 1–130.
- Fujikawa, T. (2004) Oribatid mites. In: A monitoring method development for the forest ecosystem conservation of the Shirakami-sanchi world heritage site and a forest management method development for harmonization with forest utilization in its surrounding areas. pp. 166–213. (In Japanese with English abstract.)
- Fujikawa, T., M. Fujita and J. Aoki (1993) Checklist of oribatid mites of Japan (Acari: Oribatida). J. Acarol. Soc. Jpn., 2, suppl. 1. 1–121. (In Japanese.)
- Grandjean, F. (1966) *Erogalumna zeucta* n. g., n. sp. Acarologia, 8: 475–498.
- Hammen, L. van der (1980) Glossary of acarological terminology. vol. 1 general terminology. Dr. W. Junk BV. —Publishers— The Hague, pp. 1–244.
- Matsuzaki, I. and J. Itakura (1991) Grazing of soil animals on plant soilborne disease. Nogyo-Gijutu, 46: 364–369. (In Japanese.)
- Nakamura, Y. (1988) The effect of soil management on the soil faunal makeup of a cropped andosol in central Japan. Soil Tillage Res., 12: 177–186.
- Nakamura, Y. (1989) Oribatids and enchytraeids in ecofarmed and conventionally farmed dryland grainfields of central Japan. Pedobiologia, 33: 389–398.
- Nakamura, Y. (1993) Relation between soil microorganisms and soil animals: Biological control of soil borne pathogen by collembola, oribatids and enchytraeids. Soil Microorganisms, 42: 43–59. (In Japanese.)
- Nakamura, Y. (1998) Environmental conservation and ecological studies of soil animals. Tohoku Agric. Res. Extra Issue, 11: 1–10. (In Japanese.)
- Nakamura, Y. (2000) Earthworms and potworms as keystone functional animals in pedospheres. Farming Japan, 34(5): 10–15.
- Nakamura, Y., T. Fujikawa and M. Fujita (2000) Long-term changes in the soil properties and soil macrofauna and mesofauna of an agricultural field in northern Japan during transition from chemical-intensive farming to nature farming. J. Crop Production, 3: 63–75.
- Nakamura, Y., J. Itakura and I. Matsuzaki (1991) Mycophagous meso soil animals from cropfields in Fukushima Pref. Edaphologia, 45: 49–54. (In Japanese with English abstract.)
- Nakamura, Y., H. Shiraishi and M. Hakai (2003) Earthworms and enchytraeid numbers in soybean-barley fields under till and no-till cropping systems in Japan during nine years. Mem. Fac. Agr., Ehime Univ., 48: 19–30.
- Shiraishi, H. and J. Aoki (1994) A new species of the genus *Hypochthoniella* (Acari: Oribatida) from soils on the suburban roadside in Fukushima city. Edaphologia, 52: 29–32.
- Shiraishi, H., Y. Enami & S. Okano (2003) *Folsomia hidakana* (Collembola) prevents damping off disease in cabbage and Chinese cabbage by *Rhizoctonia solani*. Pedobiologia, 47: 33–38.
- Wallwork, J. A. (1976) The Distribution and Diversity of Soil Fauna. Academic Press, London, 355 pp.

## 摘 要

中村好男・藤川徳子：環境保全型農業とササラダニ類の関係についての一考察および1新種の記載

畑地土壌のササラダニ類は3群, *Macropylina*, *Apterogasterina* および *Pterogasterina*, に区分される. 無耕起や有機物被覆条件下にあるいわゆる環境保全型農業のササラダニ類は, 耕起や化成肥料施用条件に比べ *Pterogasterina* の種の割合が多く, 環境保全型農業の生物指標の可能性が見受けられる. この *Pterogasterina* に属するカザリフリソデダニ属の新種を記載した. 本種は石垣島では有機物被覆や堆肥材料として有望なヤシ林床から採集され, 深い湾入部を持つ吻と前体部, 後体部, 翼状突起及び腹板の全表面が亀甲模様をもつ特徴により同属種から区別される. この標本は故 白石啓義 (1968–2002) 氏の採集によるものである. 同氏は農林水産省において, 土壌動物研究の立場から農業の技術改革を目指していた. その功績を讃えて *Cosomogalumna hiroyoshii* sp. nov. ヒロヨシカザリフリソデダニ (新称) と命名した.